



Opinion

Diagnosis of Asthma in Adults

Introduction

Asthma is a common respiratory condition, which often proves difficult to diagnose. In the UK, roughly one quarter of adults (aged from 20-44yrs) suffer from wheeze, 15% have wheeze and breathlessness, with roughly 7% having doctor diagnosed asthma.¹ The diagnosis of asthma requires clinicians to be aware of the common risk factors, to elicit symptoms, and to use appropriate assessment tools. There is no one gold standard test or investigation for diagnosing asthma.

Defining asthma

All definitions of asthma mention the variable presence of wheeze, breathlessness, chest tightness, and cough. Usually more than one of these is present, together with variable airflow obstruction. More recent descriptions of asthma would also involve the presence of airway hyper-responsiveness and airway inflammation.

Finding patients

Factors which increase the likelihood of asthma include co-existing atopic conditions, such as eczema and allergic rhinitis, and a family history of atopy (in particular a parent or sibling with asthma). There should be a history of variable symptoms, especially wheeze, breathlessness, chest tightness or cough which is worse in the morning, or is triggered by environmental or occupational exposure. Cold air and exercise may also cause symptoms.

Finding wheeze on auscultation, variable Peak Expiratory Flow (PEF) or obstructive FEV₁/FVC ratio on spirometry (FEV₁/FVC ratio <0.70), or a raised serum eosinophil count are also strong diagnostic pointers.² Although not diagnostic, the presence of airways obstruction is a key finding; asthma is one cause of airway obstruction, though lack of obstruction on a single reading does not exclude asthma.³

Occupation and asthma

Some patients with existing asthma will find that they come across triggers at work. In addition, there is the important and separate condition of true occupational asthma, where exposure to an inhaled sensitiser

causes an individual to develop asthma. It has been estimated that one in eight adults with a new diagnosis of asthma has an occupational cause. The importance of this diagnosis cannot be overstressed, as the patient's prognosis depends upon stopping exposure to the causative agent. Primary care is ideally placed to identify such patients,⁴ and a separate PCRS-UK opinion sheet provides more detail on this subject.⁵

The most frequently reported agents include isocyanates, flour and grain dust, colophony and fluxes, latex, animals, aldehydes and wood dust. The risk of developing occupational asthma is highest in the year after onset of occupational rhinitis.⁶

Differential diagnosis

Possible differential diagnoses can be divided into those conditions where airways obstruction is present and those where it is absent. Factors such as symptom pattern, age, smoking status, occupational and family history and other associated symptoms will help to differentiate these conditions from asthma.

Airways Obstruction Absent

Causes of chronic cough,
Hyperventilation/dysfunctional breathing
Rhinitis (allergic or otherwise),
Gastro-oesophageal reflux
Vocal cord dysfunction,
Heart Failure
Pulmonary fibrosis (see opinion sheet 40 -
<http://www.pcrs-uk.org/pubs/opinionsheets.php>)

Airways Obstruction Present

COPD,
Bronchiectasis,
Inhaled Foreign Body
Large Airways Stenosis,
Lung Cancer,
Sarcoidosis
Obliterative Bronchiolitis.

A key factor in differentiating asthma from other obstructive lung diseases is the presence or absence of reversibility. In asthma, obstruction is reversed by bronchodilators or corticosteroids, as described in Figure 2.

Making a diagnosis


Because the diagnosis of asthma is a clini-

cal one, all aspects of history, examination, investigation, and treatment trials can, and should be used when assessing patients. The presence or absence of key features in history and examination, together with lung function assessment allows clinicians to categorise patients into three levels diagnostic likelihood. These are;

- High probability of asthma
 - Key features in history and examination and abnormal lung function
- Intermediate probability of asthma
 - Some features in history and examination, inconclusive lung function
- Low probability of asthma
 - No features in history and examination, other diagnosis likely

The diagnostic process might be described as follows;

Figure 1: The diagnostic process



Lung function

In asthma, airways obstruction is, by definition variable, and reversible, and can commonly be measured in primary care using two methods, PEF monitoring and spirometry. It is clear from the current UK guidelines that each has a part to play, which can be summarised as follows.

FEV₁/FVC ratio


Spirometric measurement of FEV₁/FVC Ratio by a trained operator is an accurate and reproducible test. Patients with a ratio of <0.7 are defined as having airways obstruction, though the ratio falls with age and a slightly reduced ratio may be normal

in the elderly. A patient with asthma may, however, have a normal ratio, especially when asymptomatic. When this happens, other methods of assessment may be needed, such as serial lung function measurement.

Where obstruction is found, patients should be tested again following the administration of an effective dose of inhaled bronchodilator, or a trial of inhaled/oral corticosteroids. An improvement of 400mls in FEV₁ is strongly suggestive of asthma. It should be noted that such an improvement will not be possible in some people, particularly older or shorter patients. A PCRS-UK Spirometry opinion sheet is available for more information.⁷

Reversibility testing

Figure 2: Reversibility testing in patients whose FEV₁/FVC ratio is <70%



Peak expiratory flow

PEF is a useful tool for demonstrating variability of airflow obstruction. It is portable, inexpensive, and can be easily assessed and recorded by the patient. Its role in monitoring established asthma is well recognised.


Due to concerns regarding individual device accuracy and variability, confirming reversibility is best done using FEV₁. This can only be achieved, of course, when obstruction is already present on initial spirometry.

Where obstruction is not present, and in patients with symptoms and signs suggesting asthma, the value of serial PEF measurement as a tool for identifying 'high probability' patients is recognised in the most recent UK guidelines. In normal controls, the upper limit for PEF variability is 20%, anything greater increases the probability of asthma.

Treatment trials and investigations

Having established the level of probability of asthma, it should be possible to define a series of appropriate interventions as follows (see PCRS-UK Quick Guide for further detailed advice on the diagnosis of asthma based on BTS/SIGN Guideline for the Management of Asthma see http://www.pcrs-uk.org/asthmaguide/asthma_guide_home.php and also PCRS-UK nurse materials including Patient Group Direction (PGD) for reversibility testing,⁸ protocol for the use of spirometry in primary care⁹ and the spirometry opinion sheet.³

Figure 3: Assessing the probability of asthma²



Reasons for referral

It is important to emphasise that an unexpectedly poor response to a trial of treatment (even in a patient where there is a high diagnostic probability of asthma) should result in reassessment of both the patient and their condition.

Particular reasons for onward referral are;

- Diagnosis unclear
- Unexpected clinical findings (ie crackles, clubbing, cyanosis, cardiac disease)
- Unexplained restrictive spirometry
- Suspected occupational asthma
- Persistent non-variable breathlessness
- Monophonic wheeze or stridor
- Prominent systemic features (myalgia, fever, weight loss)
- Chronic sputum production
- CXR shadowing
- Marked blood eosinophilia (>1 x 10.9/l)
- Poor response to asthma treatment
- Severe asthma exacerbation

Advanced diagnostic tools

Where the diagnosis of asthma remains unclear, there is a role for more advanced investigation. In particular, those patients with an intermediate probability of asthma and who have normal lung function may require additional investigation. It is important to consider referral in this group, as a diagnosis of asthma usually results in lifelong treatment, and may have a significant social and occupational impact for the individual. Currently available and useful advanced diagnostic techniques include the assessment of airway hyper-responsiveness using a bronchoconstrictor challenge (often methacholine), and tests of eosinophilic airways inflammation using exhaled nitric oxide (eNO) or sputum eosinophilia.

These tests may have the additional advantage of predicting those patients who are likely to have a good response to inhaled corticosteroids.

Conclusion

There is no single, simple diagnostic pathway for adults with suspected asthma. Rating the likelihood asthma using a three section probability scale helps clinicians to seek confirmation of their clinical findings with both lung function testing and treatment trials. Where diagnostic doubt remains, patients should be referred for further specialised investigation before confining them to an inaccurate diagnosis, and a lifetime of (potentially) ineffective treatment.

References

1. Janson C, Anton J et al. The European Community Respiratory Health Survey: what are the main results so far? European Community Respiratory Health Survey II. *Eur Respir J* 2001;18(3):598-611.
2. British Guideline on the management of Asthma, a national clinical guideline. *Thorax* 2008;63 Suppl 4:iv1-121
3. PCRS-UK Opinion Sheet 1 – Spirometry. Kaplan A, Pinnock H. March 2010. http://www.pcrs-uk.org/resources/os1_spirometry.pdf
4. Levy ML and Nicholson PJ. Occupational asthma case finding: a role for primary care. Editorial, *British Journal of General Practice* 2005;54(507):731-33.
5. PCRS-UK Opinion Sheet 14. Occupational Lung Disease. Fishwick D, Sen D. January 2011. http://www.pcrs-uk.org/opinions/occupationallungdisease_final.pdf
6. Standards of care for occupational asthma. Fishwick D, Barber CM et al. *Thorax* 2008; 63:240-50. originally published online 28 Sep 2007
7. National Institute for Health and Clinical Excellence. Chronic obstructive pulmonary disease: management of adults with chronic obstructive pulmonary disease in primary and secondary care. <http://www.nice.org.uk/>. Last accessed on 2004 Mar 21.
8. Patient Group Direction for the supply/administration of salbutamol for reversibility testing in primary care by a designated healthcare professional. Stephanie Austin. October 2008. Last accessed 09/06/09 http://www.pcrs-uk.org/resources/reversibility_pgd_final.pdf
9. Protocol for the use of spirometry in primary care. Stephanie Austin, August 2007. Last accessed 09/06/09. http://www.pcrs-uk.org/resources/protocol01_spirometry_final.pdf